

Assignment # 3.

Hugo
SA (ou)

Q1. No! Let $B = \left\{ \underbrace{\frac{|0\rangle + i|1\rangle}{\sqrt{2}}, \frac{|0\rangle - |1\rangle}{\sqrt{2}}}_{|\psi_0\rangle}, \underbrace{|0\rangle - |1\rangle}_{|\psi_1\rangle} = |- \rangle \right\}$

Measuring the first qubit of $|\Phi^+\rangle$ in B and getting $|\psi_0\rangle$ gives us the following state vector:

$$|\tilde{\Phi}^+\rangle = \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle) = 0 \times |\psi_0\rangle + 1 \times |\psi_1\rangle.$$

However, measuring the last qubit, we are guaranteed to obtain $|\psi_1\rangle$ and not $|\psi_0\rangle$.

Thus, the statement is false.

By the symmetry of the state vector, measuring the 2nd qubit and getting $|\psi_0\rangle$ also gives $|\psi_1\rangle$ as a state vector.

Q2 Define $|\psi_0\rangle = \alpha|0\rangle + \beta|1\rangle$. We measure the first qubit of the state vector $|\Phi^-\rangle$ and get the following state vector:

$$|\tilde{\Phi}^-\rangle = \frac{1}{\sqrt{2}} (\bar{\alpha}|1\rangle - \bar{\beta}|0\rangle).$$

Measuring the other qubit, we have the following probability of getting $|\psi_0\rangle$:

$$\begin{aligned} \Pr[\tilde{\Phi}^- \text{ measured as } |b_0\rangle] &= |\langle b_0 | \tilde{\Phi}^- \rangle|^2 \\ &= \left| \frac{\bar{\alpha}\bar{\beta} - \bar{\alpha}\bar{\beta}}{\sqrt{2}} \right|^2 \\ &= 0. \end{aligned}$$

Thus, we always measure $|b_1\rangle$ at the second measurement.

Conversely, by the symmetry of the system (except a “-” sign, which won't add any problem as probability of measurement uses 1-1), we get the same result if we measure the 2nd qubit and get a $|b_1\rangle$, and then measure the 1st qubit.

END of ASSIGNMENT #3