




Modelling Concurrent Systems


Exercise 1.



Q2. $TS(A_{Inc}) \parallel TS(A_{Dec})$



TS($A_{Inc} \parallel A_{Dec}$):




Q3. Les effets préservent l'invariant :

- pour $x := x+1$ si $x < 200$ et $0 \leq x \leq 200$ alors $0 \leq x+1 \leq 200$
- pour $x := x-1$ si $x > 0$ et $0 \leq x \leq 200$ alors $0 \leq x-1 \leq 200$
- pour $x := 0$ si $x = 200$ et $0 \leq x \leq 200$ alors $0 \leq 0 \leq 200$

D'où l'invariant est invariant.


Q4.



Kaboom! on a cassé l'invariant!

Exercice 2.

Q1.



Exercise 3.

we have a state $x = 1$
 $x' = 0$

$$TS(PG_1 \parallel PG_2),$$

Linear Time Properties (and a bit of Modelling)

I. Safety and Invariance

Exercise 1. Invariance is safe.

Define $P_{\text{bad}} := \{ \sigma \in (2^{AP})^* \mid \exists i. \sigma(i) \not\models \varphi \}$.

Then,

$$\sigma \in P \Leftrightarrow \forall i. \sigma(i) \models \varphi$$

$$\Leftrightarrow \forall \hat{\sigma} \subseteq_{\text{fin}} \sigma, \forall i \leq \text{length } \hat{\sigma}, \hat{\sigma}(i) \models \varphi$$

$$\Leftrightarrow \forall \hat{\sigma} \subseteq_{\text{fin}} \sigma, \text{non}(\exists i. \hat{\sigma}(i) \not\models \varphi)$$

$$\Leftrightarrow \forall \hat{\sigma} \subseteq_{\text{fin}} \sigma, \hat{\sigma} \notin P_{\text{bad}}.$$

Thus P is a safety property.

Exercise 2

1. $\emptyset$ inv. safety

2. $\{ \sigma \mid \forall i, \sigma(i) \models x=0 \}$ inv. safety

3. $\{ \sigma \mid \forall i. \sigma(i) \models \neg(x=0) \wedge \neg(x>1) \} = \{ \emptyset \}^\omega$ inv. safety

4. $\{ \sigma \mid \sigma(0) \models x=0 \}$ safety $P_{\text{bad}} = \{ \hat{\sigma} \mid \hat{\sigma}(0) \not\models x=0 \}$.

5. $\{ \sigma \mid \sigma(0) \models \neg(x=0) \}$ safety $P_{\text{bad}} := \{ \hat{\sigma} \mid \hat{\sigma}(0) \models x=0 \}$

6. $\{ \sigma \mid \sigma(0) \models x=0 \text{ and } \exists i. \sigma(i) \models x>1 \}$ neither

7. $\{ \sigma \mid \exists N \forall i \geq N, \sigma(i) \models \neg(x>1) \}$ neither

8. $\{ \sigma \mid \forall N \exists i \geq N, \sigma(i) \models x>1 \}$ neither

9. $(2^{AP})^*$ inv. safety

II Operations on Safety Properties

Exercise 3. Characterization of safety properties

Q1. We have that

$$\begin{aligned} P \text{ safety property} &\Leftrightarrow \exists P_{\text{bad}}, \forall \sigma, (\sigma \in P \Leftrightarrow \text{Pref } \sigma \cap P_{\text{bad}} = \emptyset) \\ &\Leftrightarrow \exists P_{\text{bad}}, \forall \sigma, (\sigma \in P^c \Leftrightarrow \text{Pref } \sigma \cap P_{\text{bad}} \neq \emptyset) \\ (*) &\Leftrightarrow \forall \sigma, (\sigma \in P^c \Leftrightarrow \text{Pref } \sigma \cap P = \emptyset) \end{aligned}$$

For (*), " $\Leftarrow$ " we take $P_{\text{bad}} := (Q^{AP})^* \setminus P$ and we have the required property.

" $\Rightarrow$ " We have $\sigma \in \text{Pref } \sigma$ and we can conclude.

Q2. We show $P \text{ safety} \Leftrightarrow P \supseteq \text{cl } P$.

We always have $\text{cl } P \supseteq P$.

" $\Rightarrow$ " Let $\sigma \in P^c$, we will show $\sigma \notin \text{cl } P$.
i.e. $\text{pref } \sigma \not\subseteq \text{pref } P$

we have $\hat{\sigma} \in \text{pref } \sigma$ and $\hat{\sigma} \notin \text{pref } P$ by (1)

" $\Leftarrow$ " Let $\sigma \in P^c$. we will show $\exists \hat{\sigma} \in \text{pref } \sigma$ such that $\hat{\sigma} \notin \text{pref } P$
i.e. $\text{pref } \sigma \not\subseteq \text{pref } P$.

Exercise 4. Union & intersections.

Q1. For $\sigma \in (P \cup Q)^c = P^c \cap Q^c$, there exists $\hat{\sigma}_P \subseteq_{\text{fin}} \sigma$ st $\hat{\sigma}_P \cdot (Q^{AP})^\omega \cap P = \emptyset$
and $\hat{\sigma}_Q \subseteq_{\text{fin}} \sigma$ st $\hat{\sigma}_Q \cdot (Q^{AP})^\omega \cap Q = \emptyset$.

Let $\hat{\sigma} \leq \hat{\sigma}_P$ and $\hat{\sigma} \leq \hat{\sigma}_Q$. We have

$$\begin{aligned} \hat{\sigma} \cdot (Q^{AP})^\omega \cap (P \cup Q) &= (\hat{\sigma} \cdot (Q^{AP})^\omega \cap P) \cup (\hat{\sigma} \cdot (Q^{AP})^\omega \cap Q) \\ &= (\hat{\sigma}_P \cdot (Q^{AP})^\omega \cap P) \cup (\hat{\sigma}_Q \cdot (Q^{AP})^\omega \cap Q) \\ &= \emptyset \end{aligned}$$

Thus $P \cup Q$ is a safety property.

$$Q2. (P \cap Q)_{bad} := P_{bad} \cup Q_{bad}$$

$$\sigma \in P \cap Q \Leftrightarrow \forall \hat{\sigma} \in_{fini} \sigma \quad \hat{\sigma} \notin P_{bad} \text{ and } \hat{\sigma} \notin Q_{bad}$$

$$\Leftrightarrow \forall \hat{\sigma} \in_{fini} \sigma \quad \hat{\sigma} \notin (P \cap Q)_{bad}$$

III Safety properties and Transition Systems.

Exercise 5. Finite traces

$$\begin{aligned} \text{If } TS \models P \text{ then, } & Tr_{fin}(TS) \cap P_{bad} \\ &= pref(Tr^{\omega}(TS)) \cap P_{bad} \end{aligned}$$

TD n° 3

Safety and Liveness Properties

I. Liveness properties.

Exercise 1. (Closure of liveness properties)

$$cl(P) = (2^{AP})^\omega \Leftrightarrow \forall \sigma \in (2^{AP})^\omega \quad \text{pref } \sigma \subseteq \text{pref } P$$

$$\Leftrightarrow \forall \hat{\sigma} \in (2^{AP})^* \quad \hat{\sigma} \in \text{pref } P$$

$$\Leftrightarrow \forall \hat{\sigma} \in (2^{AP})^* \quad \exists \sigma \in P, \hat{\sigma} \subseteq \sigma$$

$$\Leftrightarrow P \text{ liveness property}$$

Exercise 2 (Unions & intersections).

We proved that $cl(P \cup Q) = cl(P) \cup cl(Q)$
and $cl(P \cap Q) = cl(P) \cap cl(Q)$

thus $P \cup Q$ and $P \cap Q$ are liveness properties.

Even better: if P or Q is liveness, then $P \cup Q$ is liveness.

II Topology on infinite words

Exercise 3 (Σ^ω as a topological space)

We need to prove that $\Omega\Sigma^\omega$ is stable under arbitrary unions and finite intersections.

1) Let $\mathcal{U} \subseteq \mathcal{P}(\Sigma^*)$. Define $\bar{\mathcal{U}} := \bigcup \mathcal{U}$. We have that

$$\text{ext}(\bar{\mathcal{U}}) = \bigcup_{u \in \bar{\mathcal{U}}} \text{ext}(u) = \bigcup_{u \in \mathcal{U}} \bigcup_{u \in \mathcal{U}} \text{ext}(u)$$

Thus $\Omega\Sigma^\omega$ is stable under arbitrary unions.

2) Let $U, V \subseteq \Sigma^*$.

We have that

$$\text{ext}(U) \cap \text{ext}(V) = \bigcup_{u \in U} \bigcup_{v \in V} \text{ext}(u) \cap \text{ext}(v)$$

For some $u \in U, v \in V$, we have the 3 following cases:

a) $u \not\leq v$ and $v \not\leq u$ thus $\text{ext}(u) \cap \text{ext}(v) = \emptyset$

b) $u \leq v$ thus $\text{ext}(u) \cap \text{ext}(v) = \text{ext}(u)$

c) $v \leq u$ thus $\text{ext}(u) \cap \text{ext}(v) = \text{ext}(v)$

Define

$$W := \{u \in U \mid \exists v \in V, u \leq v\} \cup \{v \in V \mid \exists u \in U, v \leq u\}$$

and we have $\text{ext}(W) = \text{ext}(U) \cap \text{ext}(V)$.

By induction, $\Omega\Sigma^\omega$ is stable under finite intersections.

Exercise 4. (Open sets)

Let $P \subseteq \Sigma^\omega$.

P is open iff $\exists U \subseteq \Sigma^*$ $P = \bigcup_{u \in U} \text{ext}(u)$

iff $\forall \sigma \in P, \exists m \in \Sigma^*, \sigma \in \text{ext}(m)$
 $\forall m \in \Sigma^*, \text{ext}(m) \subseteq P$

iff $\forall \sigma \in P, \exists m \in \Sigma^*, \sigma \in \text{ext}(m) \subseteq P$

iff $\forall \sigma \in P, \exists \hat{\sigma} \in \Sigma^*, \hat{\sigma} \leq \sigma$ and $\text{ext}(\hat{\sigma}) \subseteq P$.

Exercise 5. (Density and liveness)

P is dense iff for any non-empty open set U' , $P \cap U' \neq \emptyset$
iff for any non-empty $U \subseteq \Sigma^*$, $P \cap \text{ext}(U) \neq \emptyset$
iff for any $u \in \Sigma^*$, $P \cap \text{ext}(uP) \neq \emptyset$
iff for any $\hat{v} \in \Sigma^*$, $\exists \sigma \in P$, $\hat{v} \subseteq \sigma$.
iff P is a liveness.

III. Decomposition Theorem.

Exercise 6. Let $P \subseteq (2^{AP})^\omega$.

Define $P_{\text{safe}} := cl(P)$ which is a safety property as $cl(P_{\text{safe}}) = cl(P) = cl(P)$
and $P_{\text{live}} = P \cup P_{\text{safe}}^C$ which is a liveness property as P_{live} is
topologically dense: if $U \subseteq \Sigma^*$, and $\text{ext}(U) \cap P = \emptyset$ then $P \subseteq \text{ext}(U)^C$
and thus $cl(P) \subseteq \text{ext}(U)^C$ so we can conclude $\text{ext}(U) \subseteq cl(P)^C$. (closed)

Thus proving the decomposition theorem.

Exercise 7.

Q1. $cl(P_1) = P_1$ thus liveness and $P_{\text{safe}} = \Sigma^\omega$

Q2. $cl(P_2) = \{\sigma \mid \sigma(0) = a\}$ thus not liveness and not safety $P_{\text{liveness}} = \Sigma^\omega \setminus \{a^\omega\}$.

Q3. $cl(P_3) = \{a^\omega\} = P_3$ thus liveness and $P_{\text{safe}} = \Sigma^\omega$

Q4. $cl(P_4) = \{\sigma \mid \sigma \text{ contains } \leq 1 \text{ b's}\} = P_4 \cup \{a^\omega\}$ thus not liveness and not safety $P_{\text{live}} = \Sigma^\omega \setminus \{a^\omega\}$.

Q5. $cl(P_5) = \Sigma^\omega$ thus liveness and $P_{\text{safe}} = \Sigma^\omega$

Q6. $cl(P_6) = \Sigma^\omega$ thus liveness and $P_{safe} = \Sigma^\omega$

Q7. $cl(P_7) = \Sigma^\omega$ thus liveness and $P_{safe} = \Sigma^\omega$

Q8. $cl(P_8) = \Sigma^\omega$ thus liveness and $P_{safe} = \Sigma^\omega$.

10 m h

Topology.

I. Topology on ω -words: Examples.

c.f. exercise 7/td3

safety $\longleftrightarrow$ closed

liveness $\longleftrightarrow$ dense

closure $\longleftrightarrow$ closure (c.f. 105)

open: 1,
 $P_1 = \text{ext}(b)$

II General Properties of Topological Spaces

Exercise 2 (Closed sets)

Q1. $\emptyset = X^c$ and $X = \emptyset^c$

Q2. C_i^c is open thus $\bigcup C_i^c$ is open and then $(\bigcup C_i^c)^c = \bigcap C_i$ is closed.

Q3. C_i^c is open thus $\bigcap C_i^c$ is open and then $(\bigcap C_i^c)^c = \bigcup C_i$ is closed.

Exercise 3 (Closed and open sets).

Q1. " $\Rightarrow$ " Take $U := A \in \Omega X$. For all $x \in A$, we have $x \in U$ and $U \subseteq A$.
" $\Leftarrow$ "

We have $A = \bigcup_{a \in A} U_a \in \Omega X$ thus A is open.

Q2.

A closed iff A^c open

iff $\forall x \in A^c, \exists U \in \Omega X, x \in U \wedge U \subseteq A^c$

iff $\forall x \notin A, \exists U \in \Omega X, x \in U \wedge U \cap A = \emptyset$

Exercise 4. (closure)

Q1. Let us show $x \notin \bar{A}$ iff $\exists N \in \mathcal{D}_x, N \cap A = \emptyset$.

" $\Rightarrow$ " Consider $x \in \bar{A}$, thus there exists a closed set $A \subseteq C \subseteq X$ such that $x \notin C$.
Take $N := C^c$ which is open and contains x . And, $A \cap N = \emptyset$.

" $\Leftarrow$ " Consider x such that there exists $N \in \mathcal{D}_x$, with $N \cap A = \emptyset$.
Then, there exists an open set $U \in \Omega_X$ such that $x \in U \subseteq N$.
Take $C := U^c$ which is closed, and $U \cap A = \emptyset$ thus $A \subseteq N$.
Also, $x \notin C$.

Q2. $\bar{A}$ is the smallest closed set containing A as closed sets are stable under arbitrary intersections and $\bar{A} := \inf \{ C \subseteq X \mid C \text{ closed \& } A \subseteq C \}$.
 $\hookrightarrow$ for $\subseteq$ set inclusion

$\bar{A} = A \Leftrightarrow$ the smallest closed set containing A is A
 $\Rightarrow A$ is closed.

Q3. A dense iff $\forall U \in \Omega_X \setminus \{\emptyset\}, A \cap U \neq \emptyset$

iff $\forall C$ closed and $C \neq X, A \not\subseteq C$

iff $\bar{A} = X$

III Topology on ω -words: Properties

Exercise 5.

Suppose $v \in u$, thus $u = vw$. Take $u \nabla \in \text{ext}(u)$ and we have $vw \nabla \in \text{ext}(v)$.

Suppose $\text{ext}(u) \subseteq \text{ext}(v)$. We have $ua^\omega, ub^\omega \in \text{ext}(u)$ thus $ua^\omega, ub^\omega \in \text{ext}(v)$.
And $v \in \text{pref}(ua^\omega) \cap \text{pref}(ub^\omega) = \text{pref } u$ thus $v \subseteq u$.

Exercise 6.

P is closed iff $\forall x \notin P, \exists V \in \Omega X, x \in V$ and $V \cap P = \emptyset$

iff $\forall x \notin P, \exists U \subseteq \Sigma^*, x \in \text{ext}(U)$ and $\text{ext}(U) \cap P = \emptyset$

iff $\forall x \notin P, \exists \hat{\sigma} \in \Sigma^*, \hat{\sigma} \leq x$ and $\text{ext}(\hat{\sigma}) \cap P = \emptyset$.

IV Bases and Subbases.

Exercise 7.

Let $(V_i)_{i \in I}$ be a family of elements of ΩX .

We write $V_i := \bigcup_{j \in I_j} U_{ij}$ where $U_{ij} \in \mathcal{B}$.

Take $\bigcup V_i = \bigcup_{\substack{i \in I \\ j \in I_j}} U_{ij} \in \Omega X$.

Let $A = \bigcup_{i \in I} A_i$ $A_i \in \mathcal{B}$ and $B = \bigcup_{j \in J} B_j$.

$A \cap B = \bigcup_{\substack{i \in I \\ j \in J}} \underbrace{(A_i \cap B_j)}_{\text{an element of } \mathcal{B}} \in \Omega X$.

By induction, ΩX is closed under finite $\cap$'s.

V. Metric Spaces

Exercise 8 (Open ball topology)

Q1. Take $A, B \in \mathcal{U}$. For every $a \in A$, there exists $\varepsilon_A^a > 0$ such that $B_{\varepsilon_A^a}(a) \subseteq A$.

For every $b \in B$, there exists $\varepsilon_B^b > 0$ such that $B_{\varepsilon_B^b}(b) \subseteq B$.

For every $x \in A \cap B$, take $\varepsilon^x := \min(\varepsilon_A^a, \varepsilon_B^b) > 0$.

We have, $B_{\varepsilon^x}(x) \subseteq A \cap B$.

Consider $(A_i)_{i \in I}$ a family of elements of ΩX .

For every $i \in I$ and $a \in A_i$, there exists $\varepsilon_a^i > 0$ such that $B_{\varepsilon_a^i}(a) \subseteq A_i$.

Let $a \in \bigcup_{i \in I} A_i$. There exists $i \in I$ such that $a \in A_i$. We have $B_{\varepsilon_a^i}(a) \subseteq A_i \subseteq \bigcup_{i \in I} A_i$.

Q2. We have $\bar{S} = \{x \in X \mid \forall N \in \mathcal{N}_x, N \cap S \neq \emptyset\}$.

We have

$$x \in \bar{S} \Rightarrow \forall N \in \mathcal{N}_x, N \cap S \neq \emptyset$$

$$\Rightarrow \forall \varepsilon > 0, B_\varepsilon(x) \cap S \neq \emptyset$$

$$B_\varepsilon(x) \in \mathcal{N}_x.$$

On the other hand,

$$\forall \varepsilon > 0, B_\varepsilon(x) \cap S \neq \emptyset \Rightarrow$$

$$\forall N \in \mathcal{N}_x, \begin{cases} N \subseteq B_\varepsilon(x) \text{ for some } \varepsilon > 0 \\ N \cap S \neq \emptyset \end{cases}$$

$$\Rightarrow \forall N \in \mathcal{N}_x, N \cap S \neq \emptyset$$

Exercise 9 (Distance on ω -words)

We trivially have $d(x, y) = 0 \Leftrightarrow x = y$ and $d(x, y) = d(y, x)$.

Let σ, τ, μ be three words on A^ω . We will show $d(\sigma, \mu) \leq d(\sigma, \tau) + d(\tau, \mu)$.


If $\sigma = \mu$, the result is trivial. Same thing for $\sigma = \tau$ or $\tau = \mu$.

Suppose $\tau \neq \mu \neq \sigma \neq \tau$.

Let $i := \min \{i \mid \sigma(i) \neq \mu(i)\}$, $j := \min \{j \mid \sigma(j) \neq \tau(j)\}$

and $k := \min \{k \mid \tau(k) \neq \mu(k)\}$.

Without loss of generality, let us suppose $j < k$ (if $j = k$ then $i = j = k$ and thus the result is trivially true)



We necessarily have $i > j$ as $\sigma|_j = \tau|_j = \mu|_j$
as $\tau|_k = \mu|_k$.

$$\text{Thus, } d(\sigma, \mu) = 2^{-i} \leq 2^{-j} \leq 2^{-j} + 2^{-k} \leq d(\sigma, \tau) + d(\tau, \mu).$$

We can conclude: (A^ω, d) is a metric space.

Exercise 10 (Open ball topology on ω -words).

1) Let $U \subseteq \Sigma^*$. Let us show that $\text{ext}(U)$ is open for the open ball topology.

Let $u\sigma \in \text{ext}(U)$. Take $\varepsilon := 2^{-\text{length}(u)}$.

We have $B_\varepsilon(u\sigma) = \{ \tau \mid \min \{i \mid (u\sigma)(i) \neq \tau(i)\} > \text{length}(u) \}$
 $= \text{ext}(u)$.

Thus $\text{ext}(U) \in \mathcal{U}$.

2) Let $V \in \mathcal{U}$. We will show that $V = \text{ext}(U)$ for some $U \subseteq \Sigma^*$.

Let $v \in V$, and let $\varepsilon_v > 0$ such that $B_{\varepsilon_v}(v) \subseteq V$.

Define $U := \{v_{1-\log_2(\varepsilon_v)} \mid v \in V\}$.

We have that :

$$\begin{aligned}\text{ext}(U) &= \bigcup_{v \in V} \text{ext}(v_{1-\log_2(\varepsilon_v)}) \\ &= \bigcup_{v \in V} B_{\varepsilon_v}(v) \\ &= V.\end{aligned}$$

Thus the two topologies coincide.